



## ApoA-V(bd) Monoclonal Antibody

| Catalog No         | BYab-00602  |
|--------------------|---|
| Isotype            | lgG   |
| Reactivity         | Human   |
| Applications       | ELISA   |
| Gene Name          | APOA5   |
| Protein Name       | Apolipoprotein A-V  |
| Immunogen          | Purified recombinant fragment of human ApoA-V expressed in E. Coli.   |
| Specificity        | ApoA-V(bd) Monoclonal Antibody detects endogenous levels of ApoA-V(bd) protein.   |
| Formulation        | Purified antibody in PBS containing 0.03% sodium azide.   |
| Source             | Monoclonal, Mouse   |
| Purification       | Affinity purification   |
| Dilution           | ELISA: 1/10000. Not yet tested in other applications.   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | APOA5; RAP3; Apolipoprotein A-V; Apo-AV; ApoA-V; Apolipoprotein A5;<br>Regeneration-associated protein 3  |
| Observed Band      |   |
| Cell Pathway       | Secreted . Early endosome . Late endosome . Golgi apparatus, trans-Golgi<br>network . In the presence of SORL1, internalized to early endosomes, sorted in a<br>retrograde fashion to late endosomes, from which a portion is sent to lysosomes<br>and degradation, another portion is sorted to the trans-Golgi network  |
| Tissue Specificity | Liver and plasma.   |
| Function           | caution: It is uncertain whether Met-1 or Met-4 is the initiator., disease: Defects in APOA5 are a cause of hyperlipoproteinemia type 5 [MIM:144650].<br>Hyperlipoproteinemia type 5 is characterized by increased amounts of chylomicrons and very low density lipoprotein (VLDL) and decreased low density lipoprotein (LDL) and high density lipoprotein (HDL) in the plasma after a fast.<br>Numerous conditions cause this phenotype, including insulin-dependent diabetes mellitus, contraceptive steroids, alcohol abuse, and glycogen storage disease type 1A (GSD1A) [MIM:232200]., disease: Defects in APOA5 are a cause of susceptibility to familial hypertriglyceridemia [MIM:145750]. It is a coronary heart disease risk factor. On a regular diet the patient demonstrates increased plasma |
|                    | Nanjing BYabscience technology Co.,Ltd  |

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务

ш. 24H 在线服务,欢迎咨询

|                           | VLDL. Plasma triglycerides are persistently increased, while plasma cholesterol and phospholipids are usually within normal limits.  |
|---------------------------|--|
| Background                | The protein encoded by this gene is an apolipoprotein that plays an important role in regulating the plasma triglyceride levels, a major risk factor for coronary artery disease. It is a component of high density lipoprotein and is highly similar to a rat protein that is upregulated in response to liver injury. Mutations in this gene have been associated with hypertriglyceridemia and hyperlipoproteinemia type 5. This gene is located proximal to the apolipoprotein gene cluster on chromosome 11q23. Alternatively spliced transcript variants encoding the same protein have been identified. [provided by RefSeq, Oct 2009], |
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |

**Products Images** 

Nanjing BYabscience technology Co.,Ltd