



# Golgin 45 Polyclonal Antibody

<b>Catalog No</b>	BYab-00699
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	BLZF1
<b>Protein Name</b>	Golgin-45
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human BLZF1. AA range:10-59
<b>Specificity</b>	Golgin 45 Polyclonal Antibody detects endogenous levels of Golgin 45 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	BLZF1; JEM1; Golgin-45; Basic leucine zipper nuclear factor 1; JEM-1; p45 basic leucine-zipper nuclear factor
<b>Observed Band</b>	47kD
<b>Cell Pathway</b>	Golgi apparatus membrane .; [Isoform 1]: Nucleus . Detected in the nucleus upon heterologous expression. Not detected in the cytoplasm. .; [Isoform 2]: Cytoplasm . Not detected in the nucleus. .
<b>Tissue Specificity</b>	Detected in adrenal gland (PubMed:9129147).
<b>Function</b>	caution:Because of the presence of a potential basic motif and leucine-zipper domain, PubMed:9129147 and PubMed:11056056 have thought that BLZF1 is a potential transcription factor. They found it localized in the nucleus, except isoform 2, which was cytoplasmic. However, homology at several typical position for basic or hydrophobic residues is missing.,function:Required for normal Golgi structure and for protein transport from the endoplasmic reticulum (ER) through the Golgi apparatus to the cell surface.,induction:Up-regulated by retinoids.,subunit:Interacts with GORASP2 and with the GTP-bound form of

Nanjing BYabscience technology Co.,Ltd



RAB2, but not with other Golgi Rab proteins. GORASP2 and BLZF1 form a RAB2 effector complex on medial Golgi.,tissue specificity:Ubiquitous. Also found in cell lines derived from several hematopoietic pathologies, such as T-cell leukemia, pro-B, pre-B, myeloma, and plasmacytoma cell lines,

#### Background

caution:Because of the presence of a potential basic motif and leucine-zipper domain, PubMed:9129147 and PubMed:11056056 have thought that BLZF1 is a potential transcription factor. They found it localized in the nucleus, except isoform 2, which was cytoplasmic. However, homology at several typical position for basic or hydrophobic residues is missing.,function:Required for normal Golgi structure and for protein transport from the endoplasmic reticulum (ER) through the Golgi apparatus to the cell surface.,induction:Up-regulated by retinoids.,subunit:Interacts with GORASP2 and with the GTP-bound form of RAB2, but not with other Golgi Rab proteins. GORASP2 and BLZF1 form a RAB2 effector complex on medial Golgi.,tissue specificity:Ubiquitous. Also found in cell lines derived from several hematopoietic pathologies, such as T-cell leukemia, pro-B, pre-B, myeloma, and plasmacytoma cell lines, but not in Burkitt lymphoma cells.,

#### matters needing attention

Avoid repeated freezing and thawing!


#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




## Products Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from Jurkat cells, using BLZF1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using BLZF1 antibody.

Nanjing BYabscience technology Co.,Ltd



**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658