



# NAC-1 Monoclonal Antibody

<b>Catalog No</b>	BYab-01007
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	BTB14
<b>Protein Name</b>	Nucleus accumbens-associated protein
<b>Immunogen</b>	Purified recombinant fragment of human NAC-1 expressed in E. Coli.
<b>Specificity</b>	NAC-1 Monoclonal Antibody detects endogenous levels of NAC-1 protein.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	Affinity purification
<b>Dilution</b>	WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. ELISA: 1/10000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	NACC1; BTBD14B; NAC1; Nucleus accumbens-associated protein 1; NAC-1; BTB/POZ domain-containing protein 14B
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Distribution in the cytoplasm is dependent on phosphorylation. .
<b>Tissue Specificity</b>	Overexpressed in several types of carcinomas including ovarian serous carcinomas. Expression levels positively correlate with tumor recurrence in ovarian serous carcinomas, and intense immunoreactivity in primary ovarian tumors predicts early recurrence. Up-regulated in ovarian carcinomas after chemotherapy, suggesting a role in development of chemotherapy resistance in ovarian cancer.
<b>Function</b>	function:Functions as a transcriptional repressor. Seems to function as a transcriptional corepressor in neuronal cells through recruitment of HDAC3 and HDAC4. Contributes to tumor progression, and tumor cell proliferation and survival. This may be mediated at least in part through repressing transcriptional activity of GADD45GIP1. Required for recruiting the proteasome from the nucleus to the cytoplasm and dendritic spines.,similarity:Contains 1 BTB (POZ) domain.,subcellular location:Distribution in the cytoplasm is dependent on phosphorylation.,subunit:Homooligomer; mediated by the BTB domain. Interacts

**Nanjing BYabs science technology Co.,Ltd**



with HDAC3 and HDAC4. Interacts (via BTB domain) with CUL3, PSMD7 AND RCOR1.,tissue specificity:Overexpressed in several types of carcinomas including ovarian serous carcinomas. Expression levels positively correlate with tumor recurrence in ovarian serous carcinomas, and intense immuno

## Background

This gene encodes a member of the BTB/POZ protein family. BTB/POZ proteins are involved in several cellular processes including proliferation, apoptosis and transcription regulation. The encoded protein is a transcriptional repressor that plays a role in stem cell self-renewal and pluripotency maintenance. The encoded protein also suppresses transcription of the candidate tumor suppressor Gadd45GIP1, and expression of this gene may play a role in the progression of multiple types of cancer. A pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Feb 2012],


## matters needing attention

Avoid repeated freezing and thawing!


## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis using NAC-1 Monoclonal Antibody against HEK293 (1) and NACC1-hlgGFc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded mammary cancer tissues (left) and ovarian cancer tissues (right) with DAB staining using NAC-1 Monoclonal Antibody.

**Nanjing BYabs science technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658