



CREB-1 Monoclonal Antibody(4B4)

Catalog No	BYab-01142
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF
Gene Name	CREB1
Protein Name	Cyclic AMP-responsive element-binding protein 1
Immunogen	Recombinant Protein of CREB-1
Specificity	The antibody detects endogenous CREB-1 protein.
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1:500-1000 IHC: 1:100-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CREB1; Cyclic AMP-responsive element-binding protein 1; CREB-1; cAMP-responsive element-binding protein 1
Observed Band	43kD
Cell Pathway	Nucleus .
Tissue Specificity	Eye,Placenta,Spleen,Testis,
Function	disease:A chromosomal aberration involving CREB1 is associated with angiomatoid fibrous histiocytoma (AFH) [MIM:612160]. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type.,function:This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. Transciption activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Implicated in synchronization of circadian rhythmicity.,PTM:Stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

TIS 23 精准的 WB 检测服务

24H 在线服务,欢迎咨询

and a Coloradoral
X31 7.64
WARRAND T
南京認識界

	circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.,similarit
Background	This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. This protein binds as a homodimer to the cAMP-responsive element, an octameric palindrome. The protein is phosphorylated by several protein kinases, and induces transcription of genes in response to hormonal stimulation of the cAMP pathway. Alternate splicing of this gene results in several transcript variants encoding different isoforms. [provided by RefSeq, Mar 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Products Images



Nanjing BYabscience technology Co.,Ltd



kDa

180

130 100 -

_

70 -

55 -

40 -

35 .

25 -

Aast


CREB1

国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Various cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-CREB1 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the Lane 1: HEK293 Lane 1: A431 antibody. Predicted band size: 40,42kDa Observed band size: 40,42kDa

Whole cell lysates of LN-18 were separated by 10% SDS-PAGE, and the membrane was blotted with anti-CREB1 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: LN-18 Predicted band size: antibody. Làne 1: LN-18 Pred 40kDa Observed band size: 38kDa



Nanjing BYabscience technology Co., Ltd




国内优质抗体供应商

海内仍须加冲供应商 —— 精准的 WB 检测服务 。

24H 在线服务, 欢迎咨询





Various whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-CREB1 antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: MCF7 Lane 2: A431 Predicted band size: 37kDa Observed band size: 45kDa

Nanjing BYabscience technology Co.,Ltd