



# Ikaros 3 Polyclonal Antibody

<b>Catalog No</b>	BYab-01823
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	IKZF3
<b>Protein Name</b>	Zinc finger protein Aiolos
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human IKZF3. AA range:361-410
<b>Specificity</b>	Ikaros 3 Polyclonal Antibody detects endogenous levels of Ikaros 3 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	IKZF3; ZNFN1A3; Zinc finger protein Aiolos; Ikaros family zinc finger protein 3
<b>Observed Band</b>	62kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm .; [Isoform 1]: Nucleus .; [Isoform 3]: Nucleus .; [Isoform 11]: Nucleus .; [Isoform 14]: Nucleus . Cytoplasm .; [Isoform 12]: Cytoplasm .
<b>Tissue Specificity</b>	Expressed most strongly in peripheral blood leukocytes, the spleen, and the thymus.
<b>Function</b>	function:Transcription factor that plays an important role in the regulation of lymphocyte differentiation.,similarity:Belongs to the Ikaros C2H2-type zinc-finger protein family.,similarity:Contains 6 C2H2-type zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5.,tissue specificity:Expressed most strongly in peripheral blood leukocytes, the spleen, and the thymus.,
<b>Background</b>	This gene encodes a member of the Ikaros family of zinc-finger proteins. Three members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This gene product is a transcription factor that is important in the regulation of B lymphocyte proliferation and differentiation. Both

Nanjing BYabscience technology Co.,Ltd



Ikaros and Aiolos can participate in chromatin remodeling. Regulation of gene expression in B lymphocytes by Aiolos is complex as it appears to require the sequential formation of Ikaros homodimers, Ikaros/Aiolos heterodimers, and Aiolos homodimers. Several alternative transcripts encoding different isoforms have been described, as well as some non-protein coding variants. [provided by RefSeq, Apr 2012],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western Blot analysis of various cells using Ikaros 3 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventibiotech, MN, USA).



Western blot analysis of lysates from COLO cells, using IKZF3 Antibody. The lane on the right is blocked with the synthesized peptide.