



# HAUSP/USP7 mouse mAb

<b>Catalog No</b>	BYab-02357
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human
<b>Applications</b>	WB
<b>Gene Name</b>	usp7
<b>Protein Name</b>	
<b>Immunogen</b>	Purified recombinant human HAUSP / USP7 protein fragments expressed in E.coli.
<b>Specificity</b>	This antibody detects endogenous levels of HAUSP / USP7 and does not cross-react with related proteins.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	wb 1:200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Deubiquitinating enzyme 7;HAUSP;herpes virus associated;Herpes virus associated ubiquitin specific protease;Herpesvirus-associated ubiquitin-specific protease;TEF 1;tef-1;TEF1;Ubiquitin carboxyl terminal hydrolase 7;Ubiquitin carboxyl-terminal hydrolase 7;Ubiquitin specific peptidase 7 (herpes virus associated);Ubiquitin specific peptidase 7;Ubiquitin specific peptidase 7 herpes virus associated;Ubiquitin specific processing protease 7;Ubiquitin specific protease 7 (herpes virus associated);Ubiquitin Specific Protease 7;Ubiquitin specific protease 7 herpes virus associated;Ubiquitin thioesterase 7;ubiquitin thiolesterase 7;Ubiquitin-specific-processing protease 7;UBP 7;UBP-7;UBP7;UBP7_HUMAN;USP 7;usp-7;USP7;VMW110-ASSOCIATED PROTEIN,135-KD.
<b>Observed Band</b>	135kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm . Nucleus, PML body . Chromosome . Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of

**Nanjing BYabscience technology Co.,Ltd**



	infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies.
<b>Tissue Specificity</b>	Expressed in neural progenitor cells (at protein level) (PubMed:21258371). Widely expressed. Overexpressed in prostate cancer.
<b>Function</b>	catalytic activity:Ubiquitin C-terminal thioester + H <sub>2</sub> O = ubiquitin + a thiol.,enzyme regulation:Inhibited by N-ethyl-maleimide (NEM) and divalent cations. Tolerates high concentrations of NaCl but is inhibited at concentrations of 195 mM and higher.,function:Cleaves ubiquitin fusion protein substrates. Deubiquitinates TP53/p53 and MDM2 and strongly stabilizes TP53 even in the presence of excess MDM2, and also induces TP53-dependent cell growth repression and apoptosis.,PTM:Polyneddylated.,PTM:Polyubiquitinated.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 MATH domain.,subcellular location:Present in a minority of ND10 nuclear bodies. Association with VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus.,subunit:Monomer. Interacts with TP53, MDM2 and UBXN6. Interacts with herpesvirus
<b>Background</b>	The protein encoded by this gene belongs to the peptidase C19 family, which includes ubiquitinyl hydrolases. This protein deubiquitinates target proteins such as p53 (a tumor suppressor protein) and WASH (essential for endosomal protein recycling), and regulates their activities by counteracting the opposing ubiquitin ligase activity of proteins such as HDM2 and TRIM27, involved in the respective process. Mutations in this gene have been implicated in a neurodevelopmental disorder. [provided by RefSeq, Mar 2016],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Nanjing BYabscience technology Co.,Ltd**


网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658



## Products Images



Western blot detection of HAUSP / USP7 in HeLa and K562 cell lysates using HAUSP / USP7 mouse mAb (1:200 diluted), with super ECL. Predicted band size: 135KDa. Observed band size: 135KDa.