



ACOT2 Polyclonal Antibody

Catalog No	BYab-02470
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	ACOT2
Protein Name	Acyl-coenzyme A thioesterase 2 mitochondrial
Immunogen	The antiserum was produced against synthesized peptide derived from human ACOT2. AA range:171-220
Specificity	ACOT2 Polyclonal Antibody detects endogenous levels of ACOT2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ACOT2; PTE2; PTE2A; Acyl-coenzyme A thioesterase 2; mitochondrial; Acyl-CoA thioesterase 2; Acyl-coenzyme A thioester hydrolase 2a; CTE-Ia; Long-chain acyl-CoA thioesterase 2; ZAP128
Observed Band	53kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Strongest expression in heart, liver, muscle and kidney. Weak in placenta and pancreas.
Function	catalytic activity:Palmitoyl-CoA + H(2)O = CoA + palmitate.,caution:Was originally (PubMed:10944470) thought to be peroxisomal but was later shown (PubMed:16940157) to be mitochondrial.,function:Acyl-CoA thioesterases are a group of enzymes that catalyze the hydrolysis of acyl-CoAs to the free fatty acid and coenzyme A (CoASH), providing the potential to regulate intracellular levels of acyl-CoAs, free fatty acids and CoASH. Displays high levels of activity on medium- and long chain acyl CoAs.,similarity:Belongs to the C/M/P thioester

Nanjing BYabscience technology Co.,Ltd



hydrolase family.,tissue specificity:Strongest expression in heart, liver, muscle and kidney. Weak in placenta and pancreas.,

Background

This gene encodes a member of the acyl-CoA thioesterase protein family, and is one of four acyl-CoA hydrolase genes located in a cluster on chromosome 14. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2012],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of VEC cells using ACOT2 Polyclonal Antibody diluted at 1:500




Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from Jurkat and HUVEC cells, using ACOT2 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd



Western blot analysis of the lysates from HepG2 cells using ACOT2 antibody.