



Endo G-L1 Polyclonal Antibody

Catalog No	BYab-02633
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	EXOG
Protein Name	Nuclease EXOG mitochondrial
Immunogen	The antiserum was produced against synthesized peptide derived from human ENDOGL1. AA range:171-220
Specificity	Endo G-L1 Polyclonal Antibody detects endogenous levels of Endo G-L1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EXOG; ENDOGL1; ENDOGL2; ENGL; Nuclease EXOG; mitochondrial; Endonuclease G-like 1; Endo G-like 1
Observed Band	41kD
Cell Pathway	Mitochondrion inner membrane .
Tissue Specificity	Ubiquitous.
Function	cofactor:Divalent metal cations.,function:Endo/exonuclease with nicking activity towards supercoiled DNA, a preference for single stranded DNA and 5'-3' exonuclease activity.,miscellaneous:The active site contains 1 hydrated divalent metal cation that has only 1 direct interaction with the protein; all other interactions are via water molecules.,similarity:Belongs to the DNA/RNA non-specific endonuclease family.,subunit:Homodimer.,tissue specificity:Ubiquitous.,
Background	This gene encodes an endo/exonuclease with 5''-3'; exonuclease activity. The encoded enzyme catalyzes the hydrolysis of ester linkages at the 5'; end of a nucleic acid chain. This enzyme is localized to the mitochondria

Nanjing BYabscience technology Co.,Ltd



and may play a role in programmed cell death. Alternatively spliced transcript variants have been described. A pseudogene exists on chromosome 18. [provided by RefSeq, Feb 2009],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using Endo G-L1 Polyclonal Antibody



Western blot analysis of lysates from HUVEC, HT-29, and HepG2 cells, using ENDOGL1 Antibody. The lane on the right is blocked with the synthesized peptide.