



ZNRF2 Polyclonal Antibody

Catalog No	BYab-02845
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	ZNRF2
Protein Name	E3 ubiquitin-protein ligase ZNRF2
Immunogen	The antiserum was produced against synthesized peptide derived from human ZNRF2. AA range:161-210
Specificity	ZNRF2 Polyclonal Antibody detects endogenous levels of ZNRF2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ZNRF2; RNF202; E3 ubiquitin-protein ligase ZNRF2; Protein Ells2; RING finger protein 202; Zinc/RING finger protein 2
Observed Band	
Cell Pathway	Endosome membrane ; Peripheral membrane protein . Lysosome membrane ; Peripheral membrane protein . Cell junction, synapse, presynaptic cell membrane ; Peripheral membrane protein .
Tissue Specificity	Highly expressed in the brain, with higher expression during development than in adult. Expressed also in mammary glands, testis, colon and kidney.
Function	domain:The RING-type zinc finger domain is required for E3 ligase activity.,function:May play a role in the establishment and maintenance of neuronal transmission and plasticity via its ubiquitin ligase activity. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Present in presynaptic plasma membranes in neurons.,subunit:Interacts with UBE2N.,tissue

Nanjing BYabscience technology Co.,Ltd



specificity:Highly expressed in the brain, with higher expression during development than in adult. Expressed also in mammary glands, testis, colon and kidney.,

Background

domain:The RING-type zinc finger domain is required for E3 ligase activity.,function:May play a role in the establishment and maintenance of neuronal transmission and plasticity via its ubiquitin ligase activity. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Present in presynaptic plasma membranes in neurons.,subunit:Interacts with UBE2N.,tissue specificity:Highly expressed in the brain, with higher expression during development than in adult. Expressed also in mammary glands, testis, colon and kidney.,


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of A549 cells, using ZNRF2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd