



## ERα (phospho Ser104) Polyclonal Antibody

| Catalog No         | BYab-03276   |
|--------------------|--|
| Isotype            | lgG  |
| Reactivity         | Human;Mouse;Rat  |
| Applications       | WB;IHC;IF;ELISA  |
| Gene Name          | ESR1   |
| Protein Name       | Estrogen receptor  |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human Estrogen Receptor-alpha around the phosphorylation site of Ser104. AA range:71-120   |
| Specificity        | Phospho-ERα (S104) Polyclonal Antibody detects endogenous levels of ERα protein only when phosphorylated at S104.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG   |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000 IF 1:50-200  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | ESR1; ESR; NR3A1; Estrogen receptor; ER; ER-alpha; Estradiol receptor;<br>Nuclear receptor subfamily 3 group A member 1  |
| Observed Band      |  |
| Cell Pathway       | [Isoform 1]: Nucleus . Cytoplasm . Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . A minor fraction is associated with the inner membrane.; [Isoform 3]: Nucleus. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Single-pass type I membrane protein. Associated with the inner membrane via palmitoylation (Probable). At least a subset exists as a transmembrane protein with a N-terminal extracellular domain; Nucleus. Golgi apparatus. Cell membrane. Colocalizes with ZDHHC7 and ZDHHC21 in the Golgi apparatus where most probably palmitoylation occurs. Associated with the plasma membrane when palmitoylated. |
| Tissue Specificity | Widely expressed (PubMed:10970861). Not expressed in the pituitary gland (PubMed:10970861). ; [Isoform 3]: Widely expressed, however not expressed in the pituitary gland.   |
|                    |  |

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



| Function                  | domain:Composed of three domains: a modulating N-terminal domain, a<br>DNA-binding domain and a C-terminal steroid-binding domain.,function:Nuclear<br>hormone receptor. The steroid hormones and their receptors are involved in the<br>regulation of eukaryotic gene expression and affect cellular proliferation and<br>differentiation in target tissues.,online information:Estrogen receptor<br>entry,polymorphism:Genetic variations in ESR1 are correlated with bone mineral<br>density (BMD). Low BMD is a risk factor for osteoporotic fracture. Osteoporosis is<br>characterized by reduced bone mineral density, disrutption of bone<br>microarchitecture, and the alteration of the amount and variety of non-collagenous<br>proteins in bone. Osteoporotic bones are more at risk of<br>fracture.,PTM:Glycosylated; contains N-acetylglucosamine, probably<br>O-linked.,PTM:Phosphorylated by cyclin A/CDK2. Phosphorylation probably<br>enhances transcri |
|---------------------------|--|
| Background                | This gene encodes an estrogen receptor, a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, but also play a role in other tissues such as bone. Estrogen receptors are also involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis. Alternative promoter usage and alternative splicing result in dozens of transcript variants, but the full-length nature of many of these variants has not been determined. [provided by RefSeq, Mar 2014],   |
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |
|                           |  |


Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



## **Products Images**



Western Blot analysis of various cells using Phospho-ERα (S104) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Estrogen Receptor-alpha (Phospho-Ser104) Antibody. The picture on the right is blocked with the phospho peptide.

Nanjing BYabscience technology Co.,Ltd