



HPRT Monoclonal Antibody

Catalog No	BYab-03402
Isotype	IgG
Reactivity	Human
Applications	WB;ELISA
Gene Name	HPRT1
Protein Name	Hypoxanthine-guanine phosphoribosyltransferase
Immunogen	Purified recombinant fragment of HPRT expressed in E. Coli.
Specificity	HPRT Monoclonal Antibody detects endogenous levels of HPRT protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HPRT1; HPRT; Hypoxanthine-guanine phosphoribosyltransferase; HGPRT; HGPRTase
Observed Band	
Cell Pathway	Cytoplasm.
Tissue Specificity	Brain,
Function	catalytic activity:GMP + diphosphate = guanine + 5-phospho-alpha-D-ribose 1-diphosphate.,catalytic activity:IMP + diphosphate = hypoxanthine + 5-phospho-alpha-D-ribose 1-diphosphate.,cofactor: Binds 2 magnesium ions per subunit. One of the ions does not make direct protein contacts.,disease:Defects in HPRT1 are the cause of gout [MIM:300323]; also known as HPRT-related gout or Kelley-Seegmiller syndrome. Gout is characterized by partial enzyme activity and hyperuricemia.,disease:Defects in HPRT1 are the cause of Lesch-Nyhan syndrome (LNS) [MIM:300322]. LNS is characterized by complete lack of enzymatic activity that results in hyperuricemia, choreoathetosis, mental retardation, and compulsive self-mutilation.,online information:Hypoxanthine-guanine phosphoribosyltransferase entry,pathway:Purine metabolism; IMP biosynthesis via salvage pathway; IMP


Nanjing BYabscience technology Co.,Ltd



from hypoxanthine: step 1/1.,similarity:B

Background	hypoxanthine phosphoribosyltransferase 1 (HPRT1) Homo sapiens The protein encoded by this gene is a transferase, which catalyzes conversion of hypoxanthine to inosine monophosphate and guanine to guanosine monophosphate via transfer of the 5-phosphoribosyl group from 5-phosphoribosyl 1-pyrophosphate. This enzyme plays a central role in the generation of purine nucleotides through the purine salvage pathway. Mutations in this gene result in Lesch-Nyhan syndrome or gout.[provided by RefSeq, Jun 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using HPRT Monoclonal Antibody against truncated HPRT recombinant protein.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658