



## MTHFR Monoclonal Antibody

BYab-03413IgGHuman;RatWB;IHC;IF;ELISAMTHFRPurified recombinant fragment of human MTHFR expressed in E. Coli.
Human;Rat WB;IHC;IF;ELISA MTHFR Methylenetetrahydrofolate reductase
WB;IHC;IF;ELISA MTHFR Methylenetetrahydrofolate reductase
MTHFR Methylenetetrahydrofolate reductase
Methylenetetrahydrofolate reductase
Purified recombinant fragment of human MTHFR expressed in E. Coli.
MTHFR Monoclonal Antibody detects endogenous levels of MTHFR protein.
Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50% glycerol.
Monoclonal, Mouse
Affinity purification
WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. ELISA: 1/10000 IF 1:50-200
1 mg/ml
≥90%
-20°C/1 year
MTHFR; Methylenetetrahydrofolate reductase
cytosol,synapse,
Brain,Liver,Lung,
catalytic activity:5-methyltetrahydrofolate + NAD(P)(+) = 5,10-methylenetetrahydrofolate + NAD(P)H.,cofactor:FAD.,disease:Defects in MTHFR are the cause of methylenetetrahydrofolate reductase deficiency (MTHFRD) [MIM:236250]. MTHFRD is autosomal recessive disorder with a wide range of features including homocysteinuria, homocysteinemia [MIM:603174], developmental delay, severe mental retardation, perinatal death, psychiatric disturbances, and later-onset neurodegenerative disorders.,disease:Defects in MTHFR may be a cause of susceptibility to folate-sensitive neural tube defects (folate-sensitive NTD) [MIM:601634]. The most common NTDs are open spina bifida (myelomeningocele) and anencephaly.,disease:Defects in MTHFR may be a cause of susceptibility to ischemic stroke [MIM:601367]; also known as cerebrovascular accident or cerebral infarction. A stroke is an acute neurologic event leadin

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Background	The protein encoded by this gene catalyzes the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, a co-substrate for homocysteine remethylation to methionine. Genetic variation in this gene influences susceptibility to occlusive vascular disease, neural tube defects, colon cancer and acute leukemia, and mutations in this gene are associated with methylenetetrahydrofolate reductase deficiency.[provided by RefSeq, Oct 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




## **Products Images**



Western Blot analysis using MTHFR Monoclonal Antibody against HEK293 (1) and MTHFR-hIgGFc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded lung cancer tissues with DAB staining using MTHFR Monoclonal Antibody.



Nanjing BYabscience technology Co.,Ltd