



TAB2 Monoclonal Antibody

Catalog No	BYab-03422
Isotype	IgG
Reactivity	Human
Applications	WB;FCM;ELISA
Gene Name	TAB2
Protein Name	TGF-beta-activated kinase 1 and MAP3K7-binding protein 2
Immunogen	Purified recombinant fragment of human TAB2 expressed in E. Coli.
Specificity	TAB2 Monoclonal Antibody detects endogenous levels of TAB2 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TAB2; KIAA0733; MAP3K7IP2; TGF-beta-activated kinase 1 and MAP3K7-binding protein 2; Mitogen-activated protein kinase kinase kinase 7-interacting protein 2; TAK1-binding protein 2; TAB-2; TGF-beta-activated kinase 1-binding protein 2
Observed Band	
Cell Pathway	Membrane ; Peripheral membrane protein . Endosome membrane ; Peripheral membrane protein . Lysosome membrane ; Peripheral membrane protein . Cytoplasm, cytosol . Following IL1 stimulation, translocation occurs from the membrane to cytosol (PubMed:10882101). Interaction with TRIM38 promotes translocation from cytosol to endosome and lysosome (PubMed:24434549). .
Tissue Specificity	Widely expressed. In the embryo, expressed in the ventricular trabeculae, endothelial cells of the conotruncal cushions of the outflow tract and in the endothelial cells lining the developing aortic valves.
Function	function:Adapter linking MAP3K7/TAK1 and TRAF6 and mediator of MAP3K7 activation in the IL1 signaling pathway.,PTM:Phosphorylated.,PTM:Ubiquitinated; following IL1 stimulation or TRAF6 overexpression.,similarity:Contains 1 CUE domain.,similarity:Contains 1 RanBP2-type zinc finger.,subcellular


Nanjing BYabscience technology Co.,Ltd




	location:Following IL1 stimulation, translocation occurs from the membrane to cytosol.,subunit:Interacts with MAP3K7 and TRAF6. Interacts with NCOR1 and HDAC3 to form a ternary complex.,tissue specificity:Widely expressed.,
Background	The protein encoded by this gene is an activator of MAP3K7/TAK1, which is required for for the IL-1 induced activation of nuclear factor kappaB and MAPK8/JNK. This protein forms a kinase complex with TRAF6, MAP3K7 and TAB1, and it thus serves as an adaptor that links MAP3K7 and TRAF6. This protein, along with TAB1 and MAP3K7, also participates in the signal transduction induced by TNFSF11/RANKI through the activation of the receptor activator of NF-kappaB (TNFRSF11A/RANK), which may regulate the development and function of osteoclasts. Studies of the related mouse protein indicate that it functions to protect against liver damage caused by chemical stressors. Mutations in this gene cause congenital heart defects, multiple types, 2 (CHTD2). Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis using TAB2 Monoclonal Antibody against HEK293 (1) and TAB2-hlgGfc transfected HEK293 (2) cell lysate.



Flow cytometric analysis of HL-60 cells using TAB2 Monoclonal Antibody (green) and negative control (purple).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658