



GAPDH mouse mAb

Catalog No	BYab-03453
Isotype	IgG
Reactivity	Human;Monkey;Rat;Mouse;Saccharomycescerevisiae;Pichiapastoris;Hamster
Applications	WB
Gene Name	gapdh
Protein Name	
Immunogen	Purified recombinant human GAPDH protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of GAPDH and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:5000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	38 kDa BFA-dependent ADP-ribosylation substrate;aging associated gene 9 protein;Aging-associated gene 9 protein;BARS-38;cb609;EC 1.2.1.12;G3P_HUMAN;G3PD;G3PDH;GAPD;GAPDH; GAPDH;Glyceraldehyde 3 phosphate dehydrogenase;Glyceraldehyde 3 phosphate dehydrogenase liver;Glyceraldehyde 3 phosphate dehydrogenase muscle;Glyceraldehyde-3-phosphate dehydrogenase;KNC-NDS6;MGC102544;MGC102546;MGC103190;MGC103191;MGC105239;MGC127711; MGC88685;OCAS, p38 component;OCT1 coactivator in S phase, 38-KD component;peptidyl cysteine S-nitrosylase GAPDH;Peptidyl-cysteine S-nitrosylase GAPDH;wu:fb33a10.
Observed Band	37kD
Cell Pathway	Cytoplasm, cytosol . Nucleus . Cytoplasm, perinuclear region . Membrane . Cytoplasm, cytoskeleton . Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261). .

Nanjing BYabscience technology Co.,Ltd




Tissue Specificity	Astrocytoma,Brain,Cajal-Retzius cell,Colon adenocarcinoma,Epitheliu
Function	catalytic activity:D-glyceraldehyde 3-phosphate + phosphate + NAD(+) = 3-phospho-D-glyceroyl phosphate + NADH.,function:Independent of its glycolytic activity it is also involved in membrane trafficking in the early secretory pathway.,online information:Glyceraldehyde 3-phosphate dehydrogenase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1.,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1/5.,PTM:Reversible S-nitrosylation of Cys-152 inhibits enzymatic activity and increases endogenous ADP-ribosylation, which inhibits the enzyme in a non-reversible manner. The latter modification is more likely to be a pathophysiological event associated with inhibition of gluconeogenesis.,sequence caution:Differs quite extensively.,similarity:Belongs to the glyceraldehyde-3-phosphate dehydrogenase fami
Background	glyceraldehyde-3-phosphate dehydrogenase(GAPDH) Homo sapiens This gene encodes a member of the glyceraldehyde-3-phosphate dehydrogenase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The encoded protein has additionally been identified to have uracil DNA glycosylase activity in the nucleus. Also, this protein contains a peptide that has antimicrobial activity against E. coli, P. aeruginosa, and C. albicans. Studies of a similar protein in mouse have assigned a variety of additional functions including nitrosylation of nuclear proteins, the regulation of mRNA stability, and acting as a transferri
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd



Products Images



Western blot detection of GAPDH in Hela, 3T3, C6, COS7, CHO-K1, Rat brain and Jurkat cell lysates using GAPDH mouse mAb (1:5000 diluted). Predicted band size: 36KDa. Observed band size: 36KDa.



Western blot detection of GAPDH in *S. cerevisiae* and *P. pastoris* cell lysates using GAPDH mouse mAb (1:2000 diluted). Predicted band size: 37KDa. Observed band size: 37KDa.