



# GCNT3 Polyclonal Antibody

<b>Catalog No</b>	BYab-03892
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	GCNT3
<b>Protein Name</b>	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human GCNT3. AA range:226-275
<b>Specificity</b>	GCNT3 Polyclonal Antibody detects endogenous levels of GCNT3 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	GCNT3; Beta-1; 3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 3; C2GnT-mucin type; C2GnT-M; hC2GnT-M; Core 2/core 4 beta-1,6-N-acetylglucosaminyltransferase; C2/4GnT
<b>Observed Band</b>	
<b>Cell Pathway</b>	Golgi apparatus membrane ; Single-pass type II membrane protein .
<b>Tissue Specificity</b>	Primarily expressed in mucus-secreting tissues. Expressed in colon, kidney, small intestine, trachea, and stomach, where mucin is produced.
<b>Function</b>	catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,3-N-acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta-D-glucosaminyl-1,6)-N-acetyl-D-galactosaminyl-R.,catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,4-N-acetyl-D-glucosaminyl-R = UDP + N-acetyl-beta-D-glucosaminyl-1,6-beta-D-galactosyl-1,4-N-acetyl-D-glucosaminyl-R.,function:Glycosyltransferase that can synthesize all known mucin beta 6 N-acetylglucosaminides. Mediates core 2 and core 4 O-glycan branching, 2

Nanjing BYabscience technology Co.,Ltd



important steps in mucin-type biosynthesis. Has also I-branching enzyme activity by converting linear into branched poly-N-acetyllactosaminoglycans, leading to introduce the blood group I antigen during embryonic development.,induction:By all-trans retinoic acid (ATRA), TNF-alpha and IL13. Strongly down-regulated in colorectal cancer.,online information:Core 2/core 4 beta-1,

**Background**

This gene encodes a member of the N-acetylglucosaminyltransferase family. The encoded protein is a beta-6-N-acetylglucosamine-transferase that catalyzes the formation of core 2 and core 4 O-glycans on mucin-type glycoproteins.[provided by RefSeq, Apr 2009],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Immunofluorescence analysis of A549 cells, using GCNT3 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human liver carcinoma tissue, using GCNT3 Antibody. The picture on the right is blocked with the synthesized peptide.