



STEAP1 Polyclonal Antibody

Catalog No	BYab-04227
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	STEAP1
Protein Name	Metalloreductase STEAP1
Immunogen	The antiserum was produced against synthesized peptide derived from human STEAP1. AA range:31-80
Specificity	STEAP1 Polyclonal Antibody detects endogenous levels of STEAP1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	STEAP1; PRSS24; STEAP; Metalloreductase STEAP1; Six-transmembrane epithelial antigen of prostate 1
Observed Band	36kD
Cell Pathway	Endosome membrane ; Multi-pass membrane protein .
Tissue Specificity	Ubiquitously expressed. Highly expressed in prostate tumors.
Function	cofactor:FAD.,function:Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Ubiquitously expressed. Highly expressed in prostate tumors.,
Background	This gene is predominantly expressed in prostate tissue, and is found to be upregulated in multiple cancer cell lines. The gene product is predicted to be a six-transmembrane protein, and was shown to be a cell surface antigen significantly expressed at cell-cell junctions. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd



matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of various cells using STEAP1 Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human prostate cancer. Antibody was diluted at 1:100(4^h overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of STEAP1 antibody in paraffin-embedded human prostate carcinoma tissue.

Nanjing BYabscience technology Co.,Ltd



Western blot analysis of lysate from A549, using STEAP1 antibody.