



# TRIM3 Polyclonal Antibody

<b>Catalog No</b>	BYab-04255
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	TRIM3
<b>Protein Name</b>	Tripartite motif-containing protein 3
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human TRIM3. AA range:1-50
<b>Specificity</b>	TRIM3 Polyclonal Antibody detects endogenous levels of TRIM3 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	TRIM3; BERP; RNF22; RNF97; Tripartite motif-containing protein 3; Brain-expressed RING finger protein; RING finger protein 22; RING finger protein 97
<b>Observed Band</b>	81kD
<b>Cell Pathway</b>	Cytoplasm . Early endosome . Golgi apparatus, trans-Golgi network . Cell projection, dendrite .
<b>Tissue Specificity</b>	Expressed in brain, heart, uterus and testis.
<b>Function</b>	domain:The interaction with myosin V is dependent upon its NHL repeats, which form a beta-propeller (NHL) domain containing six blades.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 filamin repeat.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 6 NHL repeats.,subunit:Associates with myosin V and alpha-actinin-4 (ACTN4).,tissue specificity:Expressed in brain, heart, uterus and testis.,

Nanjing BYabscience technology Co.,Ltd



---

**Background**

The protein encoded by this gene is a member of the tripartite motif (TRIM) family, also called the 'RING-B-box-coiled-coil' (RBCC) subgroup of RING finger proteins. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein localizes to cytoplasmic filaments. It is similar to a rat protein which is a specific partner for the tail domain of myosin V, a class of myosins which are involved in the targeted transport of organelles. The rat protein can also interact with alpha-actinin-4. Thus it is suggested that this human protein may play a role in myosin V-mediated cargo transport. Alternatively spliced transcript variants encoding the same isoform have been identified. [provided by RefSeq, Jul 2008],

---

**matters needing attention**

Avoid repeated freezing and thawing!

---


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


---




## Products Images



Western Blot analysis of COLO cells using TRIM3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunofluorescence analysis of NIH/3T3 cells, using TRIM3 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using TRIM3 Antibody. The picture on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd



Western blot analysis of lysates from COLO cells, using TRIM3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from Jurkat cells using TRIM3 antibody.