



β-actin Monoclonal Antibody(5B7), AbFluor™ 405 Conjugated

lsotype ାତ୍	3Yab-04701 IgG
	gG
Reactivity H	
	Human;Rat;Mouse;Mk;Dg;Ch;Hamster;Rabbit;Insect
Applications IF	F;WB;IHC
Gene Name A	аств
Protein Name A	Actin cytoplasmic 1
Immunogen	
Specificity β	3-actin Monoclonal Antibody(5B7) AbFluor™ 405 Conjugated specially designed for your Immunofluorescence analysis.
	Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.
Source M	Monoclonal, Mouse IgG1
	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution C	Optimal working dilutions should be determined experimentally by the nvestigator. Suggested starting dilutions are as follows: IHC: 1:200, IF 1:200.
Concentration 1	1 mg/ml
Purity ≥	≥90%
Storage Stability -2	-20°C/1 year
Synonyms A	АСТВ
Observed Band	
Cell Pathway C	Cytoplasm, cytoskeleton . Nucleus . Localized in cytoplasmic mRNP granules containing untranslated mRNAs
	B-cell lymphoma,Brain,Cajal-Retzius cell,Eye,Fetal brain cortex,Foreskin,Hepatocellular car
[N b a d lc ty	disease:Defects in ACTB are a cause of dystonia juvenile-onset (DYTJ) [MIM:607371]. DYTJ is a form of dystonia with juvenile onset. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYTJ patients manifest progressive, generalized, dopa-unresponsive dystonia, developmental malformations and sensory hearing oss.,function:Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.,miscellaneous:In vertebrates 3 main groups of actin isoforms, alpha, beta

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins coexist in most cell types as components of the cytoskeleton and as mediators of internal cell motility.,similarity:Belongs to the
Background	This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, and integrity. This actin is a major constituent of the contractile apparatus and one of the two nonmuscle cytoskeletal actins. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd