



# KAT6B Polyclonal Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | BYab-05011   |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human;Mouse  |
| <b>Applications</b>       | WB;ELISA   |
| <b>Gene Name</b>          | KAT6B KIAA0383 MORF MOZ2 MYST4   |
| <b>Protein Name</b>       | Histone acetyltransferase KAT6B (EC 2.3.1.48) (Histone acetyltransferase MOZ2) (MOZ, YBF2/SAS3, SAS2 and TIP60 protein 4) (MYST-4) (Monocytic leukemia zinc finger protein-related factor)   |
| <b>Immunogen</b>          | Synthesized peptide derived from human protein . at AA range: 1170-1250  |
| <b>Specificity</b>        | KAT6B Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Source</b>             | Polyclonal, Rabbit,IgG   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000 ELISA 1:5000-20000   |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | ≥90%   |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           |  |
| <b>Observed Band</b>      | 228kD  |
| <b>Cell Pathway</b>       | Nucleus .  |
| <b>Tissue Specificity</b> | Ubiquitously expressed, with high levels in heart, pancreas, testis and ovary.   |
| <b>Function</b>           | catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,disease:A chromosomal aberration involving MYST4 may be a cause acute myeloid leukemias. Translocation t(10;16)(q22;p13) with CREBBP.,domain:The N-terminus is involved in transcriptional activation while the C-terminus is involved in transcriptional repression.,function:Histone acetyltransferase which may be involved in both positive and negative regulation of transcription. Required for RUNX2-dependent transcriptional activation. May be involved in cerebral cortex development. Component of the MOZ/MORF complex which has a histone H3 acetyltransferase activity.,PTM:Autoacetylated.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,similarity:Contains 2 |

Nanjing BYabscience technology Co.,Ltd



PHD-type zinc fingers.,subunit:Component of the MOZ/MORF composed

**Background**

The protein encoded by this gene is a histone acetyltransferase and component of the MOZ/MORF protein complex. In addition to its acetyltransferase activity, the encoded protein has transcriptional activation activity in its N-terminal end and transcriptional repression activity in its C-terminal end. This protein is necessary for RUNX2-dependent transcriptional activation and could be involved in brain development. Mutations have been found in patients with genitopatellar syndrome. A translocation of this gene and the CREBBP gene results in acute myeloid leukemias. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

Nanjing BYabscience technology Co.,Ltd