



CHSTC Polyclonal Antibody

Catalog No	BYab-05420
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CHST12 UNQ500/PRO1017
Protein Name	Carbohydrate sulfotransferase 12 (EC 2.8.2.5) (Chondroitin 4-O-sulfotransferase 2) (Chondroitin 4-sulfotransferase 2) (C4ST-2) (C4ST2) (Sulfotransferase Hlo)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CHSTC Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	45kD
Cell Pathway	Golgi apparatus membrane ; Single-pass type II membrane protein .
Tissue Specificity	Widely expressed. Expressed a high level in spinal chord, heart, spleen, thyroid, pituitary gland, adrenal gland, peripheral blood leukocytes, thymus, lung, small intestine, fetal kidney, fetal spleen and fetal lung.
Function	catalytic activity:3'-phosphoadenylyl sulfate + chondroitin = adenosine 3',5'-bisphosphate + chondroitin 4'-sulfate.,function:Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin and desulfated dermatan sulfate. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Activity toward partially desulfated dermatan sulfate is however lower. Does not form 4, 6-di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.,similarity:Belongs to the sulfotransferase 2 family.,tissue specificity:Widely expressed. Expressed a high level in spinal chord, heart, spleen, thyroid, pituitary gland, adrenal gland, peripheral blood leukocytes,

Nanjing BYabscience technology Co.,Ltd



thymus, lung, small intestine, fetal kidney, fetal spleen and fetal lung.,

Background

The protein encoded by this gene belongs to the sulfotransferase 2 family. It is localized to the golgi membrane, and catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin and desulfated dermatan sulfate. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage, and is distributed on the surfaces of many cells and extracellular matrices. Alternatively spliced transcript variants differing only in their 5' UTRs have been found for this gene. [provided by RefSeq, Aug 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images