



# MARH9 Polyclonal Antibody

<b>Catalog No</b>	BYab-05557
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	MARCH9 RNF179
<b>Protein Name</b>	E3 ubiquitin-protein ligase MARCH9 (EC 6.3.2.-) (Membrane-associated RING finger protein 9) (Membrane-associated RING-CH protein IX) (MARCH-IX) (RING finger protein 179)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	MARH9 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	38kD
<b>Cell Pathway</b>	Golgi apparatus membrane ; Multi-pass membrane protein . Lysosome membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Ubiquitously expressed.
<b>Function</b>	domain:The RING-CH-type zinc finger domain is required for E3 ligase activity.,function:E3 ubiquitin-protein ligase that may mediate ubiquitination of MHC-I, CD4 and ICAM1, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 RING-CH-type zinc finger.,subunit:Homodimer.,tissue specificity:Ubiquitously expressed.,
<b>Background</b>	MARCH9 is a member of the MARCH family of membrane-bound E3 ubiquitin ligases (EC 6.3.2.19). MARCH enzymes add ubiquitin (see MIM 191339) to target

**Nanjing BYabscience technology Co.,Ltd**



lysines in substrate proteins, thereby signaling their vesicular transport between membrane compartments. MARCH9 induces internalization of several membrane glycoproteins and directs them to the endosomal compartment (Bartee et al., 2004 [PubMed 14722266]; Hoer et al., 2007 [PubMed 17174307]).[supplied by OMIM, Apr 2010],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night