



# IF2M Polyclonal Antibody

<b>Catalog No</b>	BYab-06317
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	MTIF2
<b>Protein Name</b>	Translation initiation factor IF-2, mitochondrial (IF-2(Mt)) (IF-2Mt) (IF2(mt))
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 450-530
<b>Specificity</b>	IF2M Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	79kD
<b>Cell Pathway</b>	Mitochondrion.
<b>Tissue Specificity</b>	Expressed in all tissues examined. Highest level in skeletal muscle.
<b>Function</b>	function:One of the essential components for the initiation of protein synthesis. Protects formylmethionyl-tRNA from spontaneous hydrolysis and promotes its binding to the 30S ribosomal subunits. Also involved in the hydrolysis of GTP during the formation of the 70S ribosomal complex.,similarity:Belongs to the IF-2 family.,subunit:Monomer.,tissue specificity:Expressed in all tissues examined. Highest level in skeletal muscle.,
<b>Background</b>	During the initiation of protein biosynthesis, initiation factor-2 (IF-2) promotes the binding of the initiator tRNA to the small subunit of the ribosome in a GTP-dependent manner. Prokaryotic IF-2 is a single polypeptide, while eukaryotic cytoplasmic IF-2 (eIF-2) is a trimeric protein. Bovine liver mitochondria contain IF-2(mt), an 85-kD monomeric protein that is equivalent to prokaryotic IF-2. The

Nanjing BYabscience technology Co.,Ltd



predicted 727-amino acid human protein contains a 29-amino acid presequence. Human IF-2(mt) shares 32 to 38% amino acid sequence identity with yeast IF-2(mt) and several prokaryotic IF-2s, with the greatest degree of conservation in the G domains of the proteins. [provided by RefSeq, Mar 2016],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**