



# VDAC3 Polyclonal Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | BYab-06393   |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human;Rat;Mouse  |
| <b>Applications</b>       | WB;ELISA   |
| <b>Gene Name</b>          | VDAC3  |
| <b>Protein Name</b>       | Voltage-dependent anion-selective channel protein 3 (VDAC-3) (hVDAC3) (Outer mitochondrial membrane protein porin 3)   |
| <b>Immunogen</b>          | Synthesized peptide derived from part region of human protein  |
| <b>Specificity</b>        | VDAC3 Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Source</b>             | Polyclonal, Rabbit,IgG   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000 ELISA 1:5000-20000   |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | ≥90%   |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           |  |
| <b>Observed Band</b>      | 31kD   |
| <b>Cell Pathway</b>       | Mitochondrion outer membrane . Membrane . May localize to non-mitochondrial membranes. .   |
| <b>Tissue Specificity</b> | Expressed in erythrocytes (at protein level) (PubMed:27641616). Widely expressed. Highest in testis (PubMed:9781040).  |
| <b>Function</b>           | domain:Consists mainly of a membrane-spanning beta-barrel formed by 19 beta-strands.,function:Forms a channel through the mitochondrial outer membrane that allows diffusion of small hydrophilic molecules.,similarity:Belongs to the eukaryotic mitochondrial porin family.,tissue specificity:Widely expressed. Highest in testis.,   |
| <b>Background</b>         | This gene encodes a voltage-dependent anion channel (VDAC), and belongs to the mitochondrial porin family. VDACS are small, integral membrane proteins that traverse the outer mitochondrial membrane and conduct ATP and other small metabolites. They are known to bind several kinases of intermediary metabolism, thought to be involved in translocation of adenine nucleotides, and are hypothesized to form part of the mitochondrial permeability transition pore, which |

**Nanjing BYabscience technology Co.,Ltd**



results in the release of cytochrome c at the onset of apoptotic cell death. Alternatively transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Oct 2011],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4°over night