



LZTR1 Polyclonal Antibody

| | |
|---------------------------|---|
| Catalog No | BYab-06536 |
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | LZTR1 TCFL2 |
| Protein Name | Leucine-zipper-like transcriptional regulator 1 (LZTR-1) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 310-390 |
| Specificity | LZTR1 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 92kD |
| Cell Pathway | Endomembrane system . Recycling endosome . Golgi apparatus . |
| Tissue Specificity | Brain,Duodenal adenocarcinoma,Fetal brain, |
| Function | developmental stage:Expressed in fetal brain, heart, kidney, liver and lung.,disease:May play a part in the etiology of the velocardiofacial/DiGeorge syndrome (VCFS/DGS), a developmental disorder characterized by structural and functional palate anomalies, conotruncal cardiac malformations, immunodeficiency, hypocalcemia, and typical facial anomalies. Most cases result from a deletion of chromosome 22q11.2 (the DiGeorge syndrome chromosome region, or DGCR).,function:Probable transcriptional regulator that may play a crucial role in embryogenesis.,similarity:Contains 2 BTB (POZ) domains.,similarity:Contains 6 Kelch repeats., |
| Background | This gene encodes a member of the BTB-kelch superfamily. Initially described as a putative transcriptional regulator based on weak homology to members of the |

Nanjing BYabscience technology Co.,Ltd



basic leucine zipper-like family, the encoded protein subsequently has been shown to localize exclusively to the Golgi network where it may help stabilize the Gogli complex. Deletion of this gene may be associated with DiGeorge syndrome. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images