



# ZNF71 Polyclonal Antibody

<b>Catalog No</b>	BYab-06578
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	ZNF71 EZFIT
<b>Protein Name</b>	Endothelial zinc finger protein induced by tumor necrosis factor alpha (Zinc finger protein 71)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	ZNF71 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	53kD
<b>Cell Pathway</b>	Nucleus .
<b>Tissue Specificity</b>	Highly expressed in placenta, followed by brain, testis, pancreas, heart, small intestine, muscle, uterus, prostate and peripheral blood leukocytes. Not detected in liver, lung, colon, stomach, salivary and thyroid gland.
<b>Function</b>	function:May be involved in transcriptional regulation.,induction:By tumor necrosis factor alpha (TNFalpha).,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 13 C2H2-type zinc fingers.,tissue specificity:Highly expressed in placenta, followed by brain, testis, pancreas, heart, small intestine, muscle, uterus, prostate and peripheral blood leukocytes. Not detected in liver, lung, colon, stomach, salivary and thyroid gland.,
<b>Background</b>	function:May be involved in transcriptional regulation.,induction:By tumor necrosis factor alpha (TNFalpha).,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 13 C2H2-type zinc fingers.,tissue specificity:Highly expressed in placenta, followed by brain, testis, pancreas, heart,

**Nanjing BYabscience technology Co.,Ltd**



small intestine, muscle, uterus, prostate and peripheral blood leukocytes. Not detected in liver, lung, colon, stomach, salivary and thyroid gland.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**