




# VMAC Polyclonal Antibody

<b>Catalog No</b>	BYab-06811
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	VMAC
<b>Protein Name</b>	Vimentin-type intermediate filament-associated coiled-coil protein
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	VMAC Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	18kD
<b>Cell Pathway</b>	Cytoplasm . Colocalizes with vimentin-type intermediate filaments. .
<b>Tissue Specificity</b>	Cervix,Thymus,
<b>Function</b>	subcellular location:Colocalizes with vimentin-type intermediate filaments.,
<b>Background</b>	subcellular location:Colocalizes with vimentin-type intermediate filaments.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd



## Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night