



# VASN Polyclonal Antibody

<b>Catalog No</b>	BYab-06905
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	VASN SLITL2 UNQ314/PRO357/PRO1282
<b>Protein Name</b>	Vasorin (Protein slit-like 2)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	VASN Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	74kD
<b>Cell Pathway</b>	Membrane ; Single-pass type I membrane protein . Secreted .
<b>Tissue Specificity</b>	Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain, heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression is found in the tunica media of the proximal ascending aorta, the descending thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney, expression is found in the interstitial cells.
<b>Function</b>	function:May act as an inhibitor of TGF-beta signaling.,PTM:N-glycosylated.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 11 LRR (leucine-rich) repeats.,subunit:Interacts with TGFB1, TGFB2 and TGFB3.,tissue specificity:Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain, heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression is found in the tunica media of the proximal ascending aorta, the descending thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney, expression is found

**Nanjing BYabscience technology Co.,Ltd**



in the interstitial cells.,

**Background**

function:May act as an inhibitor of TGF-beta signaling.,PTM:N-glycosylated.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 11 LRR (leucine-rich) repeats.,subunit:Interacts with TGFB1, TGFB2 and TGFB3.,tissue specificity:Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain, heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression is found in the tunica media of the proximal ascending aorta, the descending thoracic aorta, the abdominal aorta and the coronary arteries. Within the kidney, expression is found in the interstitial cells.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**