



# LIGO1 Polyclonal Antibody

<b>Catalog No</b>	BYab-07156
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	LINGO1 LERN1 LRRN6A UNQ201/PRO227
<b>Protein Name</b>	Leucine-rich repeat and immunoglobulin-like domain-containing nogo receptor-interacting protein 1 (Leucine-rich repeat and immunoglobulin domain-containing protein 1) (Leucine-rich repeat neuronal pro
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 370-450
<b>Specificity</b>	LIGO1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	68kD
<b>Cell Pathway</b>	Cell membrane ; Single-pass type I membrane protein .
<b>Tissue Specificity</b>	Expressed exclusively in the central nervous system. Highest level in the in amygdala, hippocampus, thalamus and cerebral cortex. In the rest of the brain a basal expression seems to be always present. Up-regulated in substantia nigra neurons from Parkinson disease patients.
<b>Function</b>	function:Functional component of the Nogo receptor signaling complex (RTN4R/NGFR) in RhoA activation responsible for some inhibition of axonal regeneration by myelin-associated factors. Is also an important negative regulator of oligodendrocyte differentiation and axonal myelination.,PTM:N-glycosylated. Contains predominantly high-mannose glycans.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 13 LRR (leucine-rich) repeats.,subunit:Homotetramer. Forms ternary complex with RTN4R/NGFR and RTN4R/TNFRSF19.,tissue specificity:Expressed exclusively in the central

Nanjing BYabscience technology Co.,Ltd



nervous system. Highest level in the in amygdala, hippocampus, thalamus and cerebral cortex. In the rest of the brain a basal expression seems to be always present. Up-regulated in substantia nigra neurons from Parkinson disease patients.,

**Background**

function:Functional component of the Nogo receptor signaling complex (RTN4R/NGFR) in RhoA activation responsible for some inhibition of axonal regeneration by myelin-associated factors. Is also an important negative regulator of oligodentocyte differentiation and axonal myelination.,PTM:N-glycosylated. Contains predominantly high-mannose glycans.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 13 LRR (leucine-rich) repeats.,subunit:Homotetramer. Forms ternary complex with RTN4R/NGFR and RTN4R/TNFRSF19.,tissue specificity:Expressed exclusively in the central nervous system. Highest level in the in amygdala, hippocampus, thalamus and cerebral cortex. In the rest of the brain a basal expression seems to be always present. Up-regulated in substantia nigra neurons from Parkinson disease patients.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**