



# NPB Polyclonal Antibody

<b>Catalog No</b>	BYab-07353
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	NPB PPL7 PPNPB
<b>Protein Name</b>	Neuropeptide B (Preproprotein L7) (hPPL7) [Cleaved into: Neuropeptide B-23 (NPB23) (hL7); Neuropeptide B-29 (NPB29) (hL7C)]
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 1-50
<b>Specificity</b>	NPB Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	13kD
<b>Cell Pathway</b>	Secreted.
<b>Tissue Specificity</b>	Widely expressed in the central nervous system. High levels are found in substantia nigra, hypothalamus, hippocampus, spinal cord, placenta and fetal brain; lower levels are found in testis, uterus and ovary. Also detected at high levels in colorectal adenocarcinoma.
<b>Function</b>	function:May be involved in the regulation of feeding, neuroendocrine system, memory, learning and in the afferent pain pathway.,similarity:Belongs to the neuropeptide B/W family.,tissue specificity:Widely expressed in the central nervous system. High levels are found in substantia nigra, hypothalamus, hippocampus, spinal cord, placenta and fetal brain; lower levels are found in testis, uterus and ovary. Also detected at high levels in colorectal adenocarcinoma.,
<b>Background</b>	This gene encodes a member of the neuropeptide B/W family of proteins and preproprotein that is proteolytically processed to generate multiple protein

**Nanjing BYabscience technology Co.,Ltd**



products. The encoded products include neuropeptide B-23 and a C-terminally extended form, neuropeptide B-29, which are characterized by an N-terminal brominated tryptophan amino acid. Both of the encoded peptides bind with higher affinity to neuropeptide B/W (NPB/W) receptor 1 compared to the related NPB/W receptor 2. These peptides may regulate feeding, pain perception, and stress in rodents. [provided by RefSeq, Jul 2015],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night