



LCAP Polyclonal Antibody

Catalog No	BYab-07655
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	LNPEP OTASE
Protein Name	Leucyl-cystinyl aminopeptidase (Cystinyl aminopeptidase) (EC 3.4.11.3) (Insulin-regulated membrane aminopeptidase) (Insulin-responsive aminopeptidase) (IRAP) (Oxytocinase) (OTase) (Placental leucine a
Immunogen	Synthesized peptide derived from part region of human protein AA range: 705-755
Specificity	LCAP Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	112kD
Cell Pathway	Cell membrane ; Single-pass type II membrane protein . In brain only the membrane-bound form is found. The protein resides in intracellular vesicles together with GLUT4 and can then translocate to the cell surface in response to insulin and/or oxytocin. Localization may be determined by dileucine internalization motifs, and/or by interaction with tankyrases.; [Leucyl-cystinyl aminopeptidase, pregnancy serum form]: Secreted. During pregnancy serum levels are low in the first trimester, rise progressively during the second and third trimester and decrease rapidly after parturition.
Tissue Specificity	Highly expressed in placenta, heart, kidney and small intestine. Detected at lower levels in neuronal cells in the brain, in skeletal muscle, spleen, liver, testes and colon.
Function	alternative products:Experimental confirmation may be lacking for some isoforms,catalytic activity:Release of an N-terminal amino acid, Cys- -Xaa-, in

Nanjing BYabscience technology Co.,Ltd



which the half-cystine residue is involved in a disulfide loop, notably in oxytocin or vasopressin. Hydrolysis rates on a range of aminoacyl arylamides exceed that for the cystinyl derivative, however. cofactor: Binds 1 zinc ion per subunit. function: Release of an N-terminal amino acid, cleaves before cysteine, leucine as well as other amino acids. Degrades peptide hormones such as oxytocin, vasopressin and angiotensin III, and plays a role in maintaining homeostasis during pregnancy. May be involved in the inactivation of neuronal peptides in the brain. Cleaves Met-enkephalin and dynorphin. Binds angiotensin IV and may be the angiotensin IV receptor in the brain. PTM: N-glycosylated. PTM: The pregnancy serum form is derived from the membrane

Background

This gene encodes a zinc-dependent aminopeptidase that cleaves vasopressin, oxytocin, lys-bradykinin, met-enkephalin, dynorphin A and other peptide hormones. The protein can be secreted in maternal serum, reside in intracellular vesicles with the insulin-responsive glucose transporter GLUT4, or form a type II integral membrane glycoprotein. The protein catalyzes the final step in the conversion of angiotensinogen to angiotensin IV (AT4) and is also a receptor for AT4. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images