



# DCAM Polyclonal Antibody

<b>Catalog No</b>	BYab-07759
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	AMD1 AMD
<b>Protein Name</b>	S-adenosylmethionine decarboxylase proenzyme (AdoMetDC) (SAMDC) (EC 4.1.1.50) [Cleaved into: S-adenosylmethionine decarboxylase alpha chain; S-adenosylmethionine decarboxylase beta chain]
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	DCAM Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	36kD
<b>Cell Pathway</b>	cytosol,
<b>Tissue Specificity</b>	Heart,Placenta,Prostate,Thymus,Trachea,
<b>Function</b>	catalytic activity:S-adenosyl-L-methionine = (5-deoxy-5-adenosyl)(3-aminopropyl)-methylsulfonium salt + CO(2).,cofactor:Pyruvoyl group.,enzyme regulation:Both proenzyme processing and catalytic activity are stimulated by putrescine. Catalytic activity is inhibited by iodoacetic acid.,pathway:Amine and polyamine biosynthesis; S-adenosylmethioninamine biosynthesis; S-adenosylmethioninamine from S-adenosyl-L-methionine: step 1/1.,PTM:Is synthesized initially as an inactive proenzyme. Formation of the active enzyme involves a self-maturation process in which the active site pyruvoyl group is generated from an internal serine residue via an autocatalytic post-translational modification. Two non-identical subunits are generated from the proenzyme in this reaction, and the pyruvate is formed at the

Nanjing BYabscience technology Co.,Ltd



N-terminus of the alpha chain, which is derived from the carboxyl end of the proenzyme. The post-

**Background**

This gene encodes an important intermediate enzyme in polyamine biosynthesis. The polyamines spermine, spermidine, and putrescine are low-molecular-weight aliphatic amines essential for cellular proliferation and tumor promotion. Multiple alternatively spliced transcript variants have been identified. Pseudogenes of this gene are found on chromosomes 5, 6, 10, X and Y. [provided by RefSeq, Dec 2013],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

