



PLCD4 rabbit pAb

| | |
|---------------------------|--|
| Catalog No | BYab-08184 |
| Isotype | IgG |
| Reactivity | Human; Mouse;Rat |
| Applications | WB |
| Gene Name | PLCD4 |
| Protein Name | PLCD4 |
| Immunogen | Synthesized peptide derived from human PLCD4 AA range: 82-132 |
| Specificity | This antibody detects endogenous levels of PLCD4 at Human/Mouse/Rat |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.299% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase delta-4 (hPLCD4) (EC 3.1.4.11) (Phosphoinositide phospholipase C-delta-4) (Phospholipase C-delta-4) (PLC-delta-4) |
| Observed Band | 85kD |
| Cell Pathway | Membrane ; Peripheral membrane protein . Nucleus . Cytoplasm . Endoplasmic reticulum . Localizes primarily to intracellular membranes mostly to the endoplasmic reticulum. |
| Tissue Specificity | Highly expressed in skeletal muscle and kidney tissues, and at moderate level in intestinal tissue. Expressed in corneal epithelial cells. |
| Function | catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Binds 3 calcium ions per subunit. Two of the calcium ions are bound to the C2 domain.,domain:The C2 domain mediates pre-localization to the membrane prior to Ca(2+) import and non-selective Ca(2+)-mediated targeting to various cellular membranes.,domain:The PDZ-binding motif mediates the interaction with GRIP1.,domain:The PH domain is not a critical determinant of the membrane localization.,function:Hydrolyzes the phosphatidylinositol 4,5-bisphosphate (PIP2) |

Nanjing BYabscience technology Co.,Ltd



to generate 2 second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3). DAG mediates the activation of protein kinase C (PKC), while IP3 releases Ca(2+) from intracellular stores. Required for acrosome reaction in sperm during fertilization, probably by acting as an

Background

This gene encodes a member of the delta class of phospholipase C enzymes. Phospholipase C enzymes play a critical role in many cellular processes by hydrolyzing phosphatidylinositol 4,5-bisphosphate into two intracellular second messengers, inositol 1,4,5-trisphosphate and diacylglycerol. Expression of this gene may be a marker for cancer. [provided by RefSeq, Jan 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HEK293 cells, primary antibody was diluted at 1:1000, 4° over night