




## CDRT1 rabbit pAb

Catalog No	BYab-08373
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CDRT1 C17orf1 C17orf1A HREP
Protein Name	CDRT1
Immunogen	Synthesized peptide derived from human CDRT1 AA range: 364-414
Specificity	This antibody detects endogenous levels of CDRT1 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	intracellular,
Tissue Specificity	Expressed in pancreas, heart and skeletal muscle.
Function	miscellaneous:CDRT1 gene is located centromeric to and partially within proximal CMT1A-REP element. CMT1A-REP is a complex binary repeat element flanking a 1.5-Mb DNA region duplicated in Charcot-Marie-Tooth disease type I (CMT1A) or deleted in Hereditary neuropathy (HNPP).,similarity:Contains 4 WD repeats.,tissue specificity:Expressed in pancreas, heart and skeletal muscle.,
Background	Members of the F-box protein family, such as FBXW10, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603034), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM, Mar 2008],

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd