



TM14A rabbit pAb

Catalog No	BYab-09052
Isotype	IgG
Reactivity	Human; Mouse
Applications	IHC;IF
Gene Name	TMEM14A C6orf73 PTD011
Protein Name	TM14A
Immunogen	Synthesized peptide derived from human TM14A AA range: 6-56
Specificity	This antibody detects endogenous levels of TM14A at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year Mitochondrion membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane .
Synonyms Observed Band	Mitochondrion membrane ; Multi-pass membrane protein . Endoplasmic reticulum
Synonyms Observed Band Cell Pathway	Mitochondrion membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane . Expressed at significantly higher levels in ovarian cancer tissues than in normal
Synonyms Observed Band Cell Pathway Tissue Specificity	Mitochondrion membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane . Expressed at significantly higher levels in ovarian cancer tissues than in normal tissues (at protein level).
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Mitochondrion membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane . Expressed at significantly higher levels in ovarian cancer tissues than in normal tissues (at protein level).
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Mitochondrion membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane. Expressed at significantly higher levels in ovarian cancer tissues than in normal tissues (at protein level). similarity:Belongs to the UPF0136 (TMEM14) family.,


Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Products Images



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd