



YN4247

## RSLBA rabbit pAb

<b>Catalog No</b>	BYab-09087
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	RASL11A
<b>Protein Name</b>	RSLBA
<b>Immunogen</b>	Synthesized peptide derived from human RSLBA AA range: 68-118
<b>Specificity</b>	This antibody detects endogenous levels of RSLBA at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus, nucleolus . Associates with rDNA transcription unit throughout the cell cycle. .
<b>Tissue Specificity</b>	Widely expressed. Down-regulated in prostate tumors compared to normal prostate tissue. High levels found in colon tumor and normal colon tissue followed by small intestine, liver, jejunum, ileum, bladder and aorta. Lowest levels observed in endothelial cells.
<b>Function</b>	developmental stage:Down-regulated during development of THP-1 monocytes into macrophages.,induction:Down-regulated by TGFB1.,similarity:Belongs to the small GTPase superfamily. Ras family.,tissue specificity:Widely expressed. Down-regulated in prostate tumors compared to normal prostate tissue. High levels found in colon tumor and normal colon tissue followed by small intestine,


**Nanjing BYabscience technology Co.,Ltd**



liver, jejunum, ileum, bladder and aorta. Lowest levels observed in endothelial cells.,

<b>Background</b>	RASL11A is a member of the small GTPase protein family with a high degree of similarity to RAS (see HRAS, MIM 190020) proteins.[supplied by OMIM, Nov 2008],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4° over night