



Tubulin α Polyclonal Antibody

Catalog No	BYab-10638
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	Tubulin alpha
Protein Name	Tubulin alpha
Immunogen	Synthetic peptide from human protein at AA range: 380-420
Specificity	The antibody detects endogenous Tubulin α
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	Alpha-tubulin 1 antibody ALS22 antibody B ALPHA 1 antibody bA408E5.3 antibody H2 ALPHA antibody Hum a tub1 antibody Hum a tub2 antibody LIS3 antibody MGC171407 antibody MGC55332 antibody TBA4A_HUMAN antibody Testis-specific alpha-tubulin antibody TUBA1 antibody TUBA1A antibody tuba1l antibody Tuba4a antibody Tubulin alpha 1 chain antibody Tubulin alpha antibody Tubulin alpha-1 chain antibody tubulin alpha-1B chain antibody Tubulin alpha-4A chain antibody Tubulin H2-alpha antibody Tubulin, alpha 1 (testis specific) antibody tubulin, alpha 1, like antibody Tubulin, alpha 4a antibody Tubulin, alpha, testis-specific antibody Tubulin, alpha-1 antibody
Observed Band	15kD
Cell Pathway	Cytoplasm, cytoskeleton.
Tissue Specificity	Expressed at a high level in fetal brain.

Nanjing BYabscience technology Co.,Ltd



Function

disease:Defects in TUBA1A are the cause of lissencephaly type 3 (LIS3) [MIM:611603]. LIS is characterized by a smooth brain surface due to the absence (agyria) or reduction (pachygyria) of surface convolutions. It is often associated with psychomotor retardation and seizures. LIS3 features include agyria or pachygyria or laminar heterotopia, severe mental retardation, motor delay, variable presence of seizures, and abnormalities of corpus callosum, hippocampus, cerebellar vermis and brainstem.,function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,PTM:Undergoes a tyrosination/detyrosination cycle, the cyclic removal and re-addition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), resp

Background

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulins. The genes encoding these microtubule constituents belong to the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes, which are highly conserved among species. This gene encodes alpha tubulin and is highly similar to the mouse and rat Tuba1 genes. Northern blotting studies have shown that the gene expression is predominantly found in morphologically differentiated neurologic cells. This gene is one of three alpha-tubulin genes in a cluster on chromosome 12q.

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd



Products Images



Western blot analysis of HELA 293T Cell Lysate using antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000