



# PDCD6 rabbit pAb

<b>Catalog No</b>	BYab-09673
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	PDCD6 ALG2
<b>Protein Name</b>	Programmed cell death protein 6 (Apoptosis-linked gene 2 protein) (Probable calcium-binding protein ALG-2)
<b>Immunogen</b>	Synthesized peptide derived from human PDCD6 AA range: 98-148
<b>Specificity</b>	This antibody detects endogenous levels of human PDCD6
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Endoplasmic reticulum membrane ; Peripheral membrane protein . Cytoplasmic vesicle, COPII-coated vesicle membrane . Cytoplasm . Nucleus . Endosome . Interaction with RBM22 induces relocalization from the cytoplasm to the nucleus (PubMed:17045351). Translocated from the cytoplasm to the nucleus after heat shock cell treatment. Accumulates in cytoplasmic vesicle-like organelles after heat shock treatment, which may represent stress granules (PubMed:21122810). In response to calcium increase, relocates from cytoplasm to COPII vesicle coat (PubMed:27716508). Localizes to endoplasmic reticulum exit site (ERES) (PubMed:27813252)..
<b>Tissue Specificity</b>	
<b>Function</b>	
<b>Background</b>	

Nanjing BYabscience technology Co.,Ltd



**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**