



# FIG4 Polyclonal Antibody

<b>Catalog No</b>	BYab-10761
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	FIG4 KIAA0274 SAC3
<b>Protein Name</b>	FIG4
<b>Immunogen</b>	Synthesized peptide derived from human FIG4. at AA range: 341-390
<b>Specificity</b>	This antibody detects endogenous levels of FIG4
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000, ELISA 1:10000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Polyphosphoinositide phosphatase (EC 3.1.3.-) (Phosphatidylinositol 3,5-bisphosphate 5-phosphatase) (SAC domain-containing protein 3)
<b>Observed Band</b>	110kD
<b>Cell Pathway</b>	Endosome membrane . Localization requires VAC14 and PIKFYVE. .
<b>Tissue Specificity</b>	
<b>Function</b>	phospholipid metabolic process, glycerophospholipid metabolic process, vacuole organization, behavior, locomotory behavior, cell death, death, organophosphate metabolic process, neuron differentiation, phosphoinositide metabolic process, pigmentation, glycerolipid metabolic process, neuron development,
<b>Background</b>	The protein encoded by this gene belongs to the SAC domain-containing protein gene family. The SAC domain, approximately 400 amino acids in length and consisting of seven conserved motifs, has been shown to possess phosphoinositide phosphatase activity. The yeast homolog, Sac1p, is involved in the regulation of various phosphoinositides, and affects diverse cellular functions such as actin cytoskeleton organization, Golgi function, and maintenance of

Nanjing BYabscience technology Co.,Ltd



vacuole morphology. Membrane-bound phosphoinositides function as signaling molecules and play a key role in vesicle trafficking in eukaryotic cells. Mutations in this gene have been associated with Charcot-Marie-Tooth disease, type 4J. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of 1,mouse-liver 2,hela 3,mouse-brain cells using primary antibody diluted at 1:1000(4°C overnight). Secondary antibody:Goat Anti-rabbit IgG IRDye 800( diluted at 1:5000, 25°C, 1 hour)