



PEN2 rabbit pAb

Catalog No	BYab-10997
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	IHC;IF
Gene Name	PSENEN PEN2 MDS033
Protein Name	PEN2
Immunogen	Synthesized peptide derived from human PEN2 AA range: 51-101
Specificity	This antibody detects endogenous levels of PEN2 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Golgi apparatus, Golgi stack membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein . Membrane ; Multi-pass membrane protein . Predominantly located in the endoplasmic reticulum and in the cis-Golgi.
Tissue Specificity	Widely expressed. Expressed in leukocytes, lung, placenta, small intestine, liver, kidney, spleen thymus, skeletal muscle, heart and brain.
Function	function:Essential subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (beta-amyloid precursor protein). Probably represents the last step of maturation of gamma-secretase, facilitating endoproteolysis of presenilin and conferring gamma-secretase activity.,similarity:Belongs to the PEN-2 family.,subcellular location:Predominantly located in the endoplasmic reticulum and in the cis-Golgi.,subunit:Component of the gamma-secretase complex, a complex composed of a presenilin homodimer

Nanjing BYabscience technology Co.,Ltd



(PSEN1 or PSEN2), nicastrin (NCSTN), APH1 (APH1A or APH1B) and PSENEN/PEN2. Such minimal complex is sufficient for secretase activity, although other components may exist.,tissue specificity:Widely expressed. Expressed in leukocytes, lung, placenta, small intestine, liver, kidney, spleen thymus, s

Background

Presenilins, which are components of the gamma-secretase protein complex, are required for intramembranous processing of some type I transmembrane proteins, such as the Notch proteins and the beta-amyloid precursor protein. Signaling by Notch receptors mediates a wide range of developmental cell fates. Processing of the beta-amyloid precursor protein generates neurotoxic amyloid beta peptides, the major component of senile plaques associated with Alzheimer's disease. This gene encodes a protein that is required for Notch pathway signaling, and for the activity and accumulation of gamma-secretase. Mutations resulting in haploinsufficiency for this gene cause familial acne inversa-2 (ACNINV2). Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).