



# MYT1L rabbit pAb

<b>Catalog No</b>	BYab-11205
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB;ELISA;IHC
<b>Gene Name</b>	MYT1L KIAA1106
<b>Protein Name</b>	MYT1L
<b>Immunogen</b>	Synthesized peptide derived from human MYT1L AA range: 595-645
<b>Specificity</b>	This antibody detects endogenous levels of MYT1L at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus . Chromosome . Preferentially binds to DNA binding sites that are in an open chromatin configuration. .
<b>Tissue Specificity</b>	
<b>Function</b>	function:May function as a panneural transcription factor associated with neuronal differentiation. May play a role in the development of neurons and oligodendrogalia in the CNS.,similarity:Contains 6 C2HC-type zinc fingers.,
<b>Background</b>	This gene encodes a member of the zinc finger superfamily of transcription factors. The encoded protein belongs to a novel class of cystein-cystein-histidine-cystein zinc finger proteins that function in the developing mammalian central nervous system. Forced expression of this gene in combination with the basic helix-loop-helix transcription factor NeuroD1 and the transcription factors POU class 3 homeobox 2 and achaete-scute family basic helix-loop-helix transcription factor 1 can convert fetal and postnatal human fibroblasts into induced neuronal cells, which are able to generate action

Nanjing BYabscience technology Co.,Ltd



potentials. Mutations in this gene have been associated with autosomal mental retardation-39 and autism spectrum disorder. Alternative splicing results in multiple variants. [provided by RefSeq, Aug 2016],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).

**Nanjing BYabscience technology Co.,Ltd**