



KBTB8 rabbit pAb

Catalog No	BYab-11207
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	KBTBD8 KIAA1842 TAKRP
Protein Name	KBTB8
Immunogen	Synthesized peptide derived from human KBTB8 AA range: 445-495
Specificity	This antibody detects endogenous levels of KBTB8 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
D!t	. 000/
Purity	≥90%
Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Cytoplasm, cytoskeleton, spindle . Golgi apparatus . Translocates to the spindle
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Cytoplasm, cytoskeleton, spindle . Golgi apparatus . Translocates to the spindle
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Cytoplasm, cytoskeleton, spindle . Golgi apparatus . Translocates to the spindle apparatus during mitosis similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Cytoplasm, cytoskeleton, spindle . Golgi apparatus . Translocates to the spindle apparatus during mitosis similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year Cytoplasm, cytoskeleton, spindle . Golgi apparatus . Translocates to the spindle apparatus during mitosis similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 Kelch repeats.,


Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd