



CB082 rabbit pAb

Catalog No	BYab-11478
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	C2orf82 UNQ830/PRO1757
Protein Name	CB082
Immunogen	Synthesized peptide derived from human CB082 AA range: 68-118
Specificity	This antibody detects endogenous levels of CB082 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band Cell Pathway	Membrane; Single-pass membrane protein. Cytoplasm. Secreted, extracellular space, extracellular matrix. Appears as reticular-like structures throughout the cytoplasm and adjacent to the plasma membrane. In proliferation and hypertrophic chondrocytes, detected intracellulary and in the pericellular extracellular matrix. In primary spongiosa, detected only in the extracellular matrix.
	space, extracellular matrix. Appears as reticular-like structures throughout the cytoplasm and adjacent to the plasma membrane. In proliferation and hypertrophic chondrocytes, detected intracellulary and in the pericellular extracellular matrix. In primary spongiosa, detected only in the extracellular
Cell Pathway	space, extracellular matrix. Appears as reticular-like structures throughout the cytoplasm and adjacent to the plasma membrane. In proliferation and hypertrophic chondrocytes, detected intracellulary and in the pericellular extracellular matrix. In primary spongiosa, detected only in the extracellular matrix
Cell Pathway Tissue Specificity	space, extracellular matrix. Appears as reticular-like structures throughout the cytoplasm and adjacent to the plasma membrane. In proliferation and hypertrophic chondrocytes, detected intracellulary and in the pericellular extracellular matrix. In primary spongiosa, detected only in the extracellular matrix
Cell Pathway Tissue Specificity Function	space, extracellular matrix. Appears as reticular-like structures throughout the cytoplasm and adjacent to the plasma membrane. In proliferation and hypertrophic chondrocytes, detected intracellulary and in the pericellular extracellular matrix. In primary spongiosa, detected only in the extracellular matrix

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd