



TRPM3 rabbit pAb

| | |
|---------------------------|---|
| Catalog No | BYab-11481 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | TRPM3 KIAA1616 LTRPC3 |
| Protein Name | TRPM3 |
| Immunogen | Synthesized peptide derived from human TRPM3 AA range: 666-716 |
| Specificity | This antibody detects endogenous levels of TRPM3 at Human |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1: 500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | Cell membrane ; Multi-pass membrane protein . |
| Tissue Specificity | Expressed primarily in the kidney and, at lower levels, in brain, testis, ovary, pancreas and spinal cord. Expression in the brain and kidney was determined at protein level. In the kidney, expressed predominantly in the collecting tubular epithelium in the medulla, medullary rays, and periglomerular regions; in the brain, highest levels are found in the cerebellum, choroid plexus, the locus coeruleus, the posterior thalamus and the substantia nigra. Down-regulated in renal tumors compared to normal kidney. |
| Function | function:Calcium channel mediating constitutive calcium ion entry. Its activity is increased by reduction in extracellular osmolarity, by store depletion and muscarinic receptor activation.,similarity:Belongs to the transient receptor family. LTrpC subfamily.,tissue specificity:Expressed primarily in the kidney and, at lower levels, in brain, testis, ovary, pancreas and spinal cord. Expression in the brain and kidney was determined at protein level. In the kidney, expressed predominantly in the collecting tubular epithelium in the medulla, medullary rays, |

Nanjing BYabscience technology Co.,Ltd



and periglomerular regions; in the brain, highest levels are found in the cerebellum, choroid plexus, the locus coeruleus, the posterior thalamus and the substantia nigra. Down-regulated in renal tumors compared to normal kidney.,

Background

The product of this gene belongs to the family of transient receptor potential (TRP) channels. TRP channels are cation-selective channels important for cellular calcium signaling and homeostasis. The protein encoded by this gene mediates calcium entry, and this entry is potentiated by calcium store depletion. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night