



## PI42C rabbit pAb

Catalog No	BYab-11899
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	PIP4K2C PIP5K2C
Protein Name	PI42C
Immunogen	Synthesized peptide derived from human PI42C AA range: 262-312
Specificity	This antibody detects endogenous levels of PI42C at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Endoplasmic reticulum . Cytoplasm .
Tissue Specificity	
Function	catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 5-phosphate = ADP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate.,function:May play an important role in the production of Phosphatidylinositol bisphosphate (PIP2), in the endoplasmic reticulum.,PTM:Phosphorylated.,similarity:Contains 1 PI5K domain.,subcellular location:Mostly found in the cytosol and surrounding plasma membrane. However, its presence in the endoplasmic reticulum seems to be a prerequisite for PIP2 synthesis.,
Background	

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd