



PRD14 rabbit pAb

Catalog No	BYab-12325
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	PRDM14
Protein Name	PRD14
Immunogen	Synthesized peptide derived from human PRD14 AA range: 391-441
Specificity	This antibody detects endogenous levels of PRD14 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in embryonic stem cells. Tends to be overexpressed in breast cancer (at protein level).
Function	function:May be involved in transcriptional regulation.,similarity:Contains 1 SET domain.,similarity:Contains 6 C2H2-type zinc fingers.,
Background	This gene encodes a member of the PRDI-BF1 and RIZ homology domain containing (PRDM) family of transcriptional regulators. The encoded protein may possess histone methyltransferase activity and plays a critical role in cell pluripotency by suppressing the expression of differentiation marker genes. Expression of this gene may play a role in breast cancer. [provided by RefSeq, Dec 2011],
matters needing attention	Avoid repeated freezing and thawing!


Nanjing BYabscience technology Co.,Ltd




Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).