



KLHL13 Monoclonal Antibody

Catalog No	BYab-12608
Isotype	IgG
Reactivity	Human
Applications	WB;IHC;IF;FCM;ELISA
Gene Name	KLHL13
Protein Name	Kelch-like protein 13
Immunogen	Purified recombinant fragment of human KLHL13 expressed in E. Coli.
Specificity	KLHL13 Monoclonal Antibody detects endogenous levels of KLHL13 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KLHL13; BKLHD2; KIAA1309; Kelch-like protein 13; BTB and kelch domain-containing protein 2
Observed Band	
Cell Pathway	midbody,Cul3-RING ubiquitin ligase complex,
Tissue Specificity	Brain,Testis,Thalamus,Whole embryo,
Function	function:Substrate-specific adapter for a CUL3-based E3 ubiquitin-protein ligase complex. Within this complex, controls the dynamic behavior of AURKB on mitotic chromosomes and thereby coordinates faithful mitotic progression and completion of cytokinesis.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 6 Kelch repeats.,subunit:Forms a complex with CUL3 and KLHL9. Interacts with AURKB. Interacts with CUL3.,
Background	This gene encodes a BTB and kelch domain containing protein and belongs to the kelch repeat domain containing superfamily of proteins. The encoded protein

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




	functions as an adaptor protein that complexes with Cullin 3 and other proteins to form the Cullin 3-based E3 ubiquitin-protein ligase complex. This complex is necessary for proper chromosome segregation and completion of cytokinesis. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Mar 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.







Products Images





Western Blot analysis using KLHL13 Monoclonal Antibody against HeLa (1) and MCF-7 (2) cell lysate.






Immunohistochemistry analysis of paraffin-embedded brain tissues (left) and pancreas tissues (right) with DAB staining using KLHL13 Monoclonal Antibody.





Immunofluorescence analysis of NTERA-2 cells (left) and U251 (right) cells using KLHL13 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of 3T3/L1 cells using KLHL13 Monoclonal Antibody (green) and negative control (purple).

Nanjing BYabscience technology Co.,Ltd