



MuSK Monoclonal Antibody

Catalog No	BYab-12931
Isotype	IgG
Reactivity	Human
Applications	IHC;IF;ELISA
Gene Name	MUSK
Protein Name	Muscle, skeletal receptor tyrosine-protein kinase
Immunogen	Purified recombinant extracellular fragment of human MuSK (aa24-209) fused with hlgGfC tag expressed in HEK293 cell line.
Specificity	MuSK Monoclonal Antibody detects endogenous levels of MuSK protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MUSK; Muscle; skeletal receptor tyrosine-protein kinase; Muscle-specific tyrosine-protein kinase receptor; MuSK; Muscle-specific kinase receptor
Observed Band	
Cell Pathway	Cell junction, synapse, postsynaptic cell membrane ; Single-pass type I membrane protein . Colocalizes with acetylcholine receptors (AChR) to the postsynaptic cell membrane of the neuromuscular junction. .
Tissue Specificity	
Function	catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in MUSK is a cause of autosomal recessive congenital myasthenic syndrome (CMS) [MIM:608931]. Congenital myasthenic syndromes are inherited disorders of neuromuscular transmission that stem from mutations in presynaptic, synaptic, or postsynaptic proteins. MUSK mutations lead to decreased agrin-dependent AChR aggregation, a critical step in the formation of the neuromuscular junction.,function:Receptor tyrosine kinase that is a key mediator of agrin's action and is involved in neuromuscular junction (NMJ) organization.,online information:MuSK entry,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Contains 1 FZ (frizzled)

Nanjing BYabscience technology Co.,Ltd



domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like) domains.,s

Background

This gene encodes a muscle-specific tyrosine kinase receptor. The encoded protein may play a role in clustering of the acetylcholine receptor in the postsynaptic neuromuscular junction. Mutations in this gene have been associated with congenital myasthenic syndrome. Alternatively spliced transcript variants have been described.[provided by RefSeq, Oct 2009],


matters needing attention

Avoid repeated freezing and thawing!



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human lung cancer (A), muscles (B) and breast cancer (C) with DAB staining using MuSK Monoclonal Antibody.



Confocal immunofluorescence analysis of HEK293 cells transfected with extracellular MUSK (aa24-209)-hlgGfc using MuSK Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658