



Cleaved-CD97 α (L530) Polyclonal Antibody

Catalog No	BYab-13776
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CD97
Protein Name	CD97 antigen
Immunogen	The antiserum was produced against synthesized peptide derived from human CD97 α . AA range:481-530
Specificity	Cleaved-CD97 α (L530) Polyclonal Antibody detects endogenous levels of fragment of activated CD97 α protein resulting from cleavage adjacent to L530.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	CD97; CD97 antigen; Leukocyte antigen CD97; CD antigen CD97
Observed Band	55kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .; [Adhesion G protein-coupled receptor E5 subunit alpha]: Secreted, extracellular space .
Tissue Specificity	Broadly expressed, found on most hematopoietic cells, including activated lymphocytes, monocytes, macrophages, dendritic cells, and granulocytes. Expressed also abundantly by smooth muscle cells. Expressed in thyroid, colorectal, gastric, esophageal and pancreatic carcinomas too. Expression are increased under inflammatory conditions in the CNS of multiple sclerosis and in synovial tissue of patients with rheumatoid arthritis. Increased expression of CD97 in the synovium is accompanied by detectable levels of soluble CD97 in the synovial fluid.
Function	domain:Binding to chondroitin sulfate is mediated by the fourth EGF domain.,domain:The first two EGF domains mediate the interaction with DAF. A third tandemly arranged EGF domain is necessary for the structural integrity of the binding region.,function:Receptor potentially involved in both adhesion and

Nanjing BYabscience technology Co.,Ltd



signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.,induction:Rapid up-regulation during lymphocyte activation.,PTM:Proteolytically cleaved into 2 subunits, an extracellular alpha subunit and a seven-transmembrane subunit.,similarity:Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS domain.,similarity:Contains 5 EGF-like domains.,subunit:Forms a heterodimer, consisting of a large extracellular region (alpha subunit) non-covalently linked to a seven-transmembrane moiety (beta subunit). Interacts

Background

This gene encodes a member of the EGF-TM7 subfamily of adhesion G protein-coupled receptors, which mediate cell-cell interactions. These proteins are cleaved by self-catalytic proteolysis into a large extracellular subunit and seven-span transmembrane subunit, which associate at the cell surface as a receptor complex. The encoded protein may play a role in cell adhesion as well as leukocyte recruitment, activation and migration, and contains multiple extracellular EGF-like repeats which mediate binding to chondroitin sulfate and the cell surface complement regulatory protein CD55. Expression of this gene may play a role in the progression of several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms with 3 to 5 EGF-like repeats have been observed for this gene. This gene is found in a cluster with other EGF-TM7 genes on the short arm of chromosome 1


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of CD97 α (Cleaved-Leu530) Antibody. The lane on the right is blocked with the CD97 α (Cleaved-Leu530) peptide.

Nanjing BYabscience technology Co.,Ltd