



# DGK- $\zeta$ Polyclonal Antibody

<b>Catalog No</b>	BYab-14725
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	DGKZ
<b>Protein Name</b>	Diacylglycerol kinase zeta
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human DGKZ. AA range:601-650
<b>Specificity</b>	DGK- $\zeta$ Polyclonal Antibody detects endogenous levels of DGK- $\zeta$ protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	DGKZ; DAGK6; Diacylglycerol kinase zeta; DAG kinase zeta; Diglyceride kinase zeta; DGK-zeta
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus . Cytoplasm, cytosol . Cell membrane . Cell projection, lamellipodium .
<b>Tissue Specificity</b>	Highest levels in brain, and substantial levels in skeletal muscle, heart, and pancreas. ; [Isoform 2]: Predominantly expressed in muscle.
<b>Function</b>	catalytic activity:ATP + 1,2-diacylglycerol = ADP + 1,2-diacyl-sn-glycerol 3-phosphate.,caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Displays a strong preference for 1,2-diacylglycerols over 1,3-diacylglycerols, but lacks substrate specificity among molecular species of long chain diacylglycerols. Isoform 2 but not isoform 1 regulates RASGRP1 activity.,PTM:Phosphorylation of the MARCKS homology domain by PKC reduces nuclear accumulation of DGK-zeta.,similarity:Belongs to the eukaryotic diacylglycerol kinase family.,similarity:Contains 1 DAGKc domain.,similarity:Contains 2 ANK repeats.,similarity:Contains 2

Nanjing BYabscience technology Co.,Ltd



phorbol-ester/DAG-type zinc fingers.,subunit:Interacts with the PDZ domain of the syntrophin SNTG1 and that of SNX27. Isoform 2 forms a signaling complex with RASGRP1 and HRAS.,tissue speci

**Background**

The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It may attenuate protein kinase C activity by regulating diacylglycerol levels in intracellular signaling cascade and signal transduction. Alternative splicing occurs at this locus and multiple transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Nov 2010],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Immunohistochemistry analysis of paraffin-embedded human brain, using DGKZ Antibody. The picture on the right is blocked with the synthesized peptide.