



# PI 3-Kinase p110 $\gamma$ Polyclonal Antibody

<b>Catalog No</b>	BYab-14914
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	PIK3CG
<b>Protein Name</b>	Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit gamma isoform
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human PIK3CG. AA range:881-930
<b>Specificity</b>	PI 3-Kinase p110 $\gamma$ Polyclonal Antibody detects endogenous levels of PI 3-Kinase p110 $\gamma$ protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	PIK3CG; Phosphatidylinositol 4; 5-bisphosphate 3-kinase catalytic subunit gamma isoform; PI3-kinase subunit gamma; PI3K-gamma; PI3Kgamma; PtdIns-3-kinase subunit gamma; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit
<b>Observed Band</b>	120kD
<b>Cell Pathway</b>	Cytoplasm . Cell membrane .
<b>Tissue Specificity</b>	Pancreas, skeletal muscle, liver and heart.
<b>Function</b>	catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate.,enzyme regulation:Activated by both the alpha and the beta-gamma G proteins.,function:3-phosphorylates the cellular phosphoinositide PtdIns-4,5-bisphosphate (PtdIns(4,5)P2) to produce PtdIns-3, 4,5-triisphosphate (PtdIns(3,4,5)P3). Links G-protein coupled receptor activation to the secondary messenger PtdIns(3,4,5)P3 production.,pathway:Phospholipid metabolism; phosphatidylinositol phosphate

Nanjing BYabscience technology Co.,Ltd



biosynthesis.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Heterodimer of a catalytic subunit (PIK3CG/p120) and a regulatory (PIK3R5a/p101) subunit.,tissue specificity:Pancreas, skeletal muscle, liver and heart.,

**Background**

Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha, p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2015],

**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




## Products Images



Western blot analysis of PIK3CG Antibody. The lane on the right is blocked with the PIK3CG peptide.



Immunohistochemistry analysis of paraffin-embedded human pancreas, using PIK3CG Antibody. The lane on the right is blocked with the PIK3CG peptide.



Western blot analysis of the lysates from Jurkat cells using PIK3CG antibody.